

FDA approves new monotherapy indication for Merck's KEYTRUDA

01 August 2019 | News

KEYTRUDA now approved for recurrent locally advanced or metastatic squamous cell carcinoma of the esophagus in patients whose tumors Express PD-L1 (CPS \geq 10) with disease progression after one or more prior lines of systemic therapy



Merck known as MSD outside the United States and Canada, has announced that the U.S. Food and Drug Administration (FDA) has approved KEYTRUDA, Merck's anti-PD-1 therapy, as monotherapy for the treatment of patients with recurrent locally advanced or metastatic squamous cell carcinoma of the esophagus whose tumors express PD-L1 (Combined Positive Score [CPS] \geq 10) as determined by an FDA-approved test, with disease progression after one or more prior lines of systemic therapy.

"Historically, patients with advanced esophageal cancer have had limited treatment options, particularly after their disease has progressed," said Dr. Jonathan Cheng, vice president, oncology clinical research, Merck Research Laboratories. "With this approval, KEYTRUDA is now the first anti-PD-1 therapy approved for the treatment of previously-treated patients with recurrent locally advanced or metastatic squamous cell carcinoma of the esophagus whose tumors express PD-L1 (CPS \geq 10), providing an important new monotherapy option for physicians and patients in the United States."

Immune-mediated adverse reactions, which may be severe or fatal, can occur with KEYTRUDA, including pneumonitis, colitis, hepatitis, endocrinopathies, nephritis and renal dysfunction, severe skin reactions, solid organ transplant rejection, and complications of allogeneic hematopoietic stem cell transplantation (HSCT). Based on the severity of the adverse reaction, KEYTRUDA should be withheld or discontinued and corticosteroids administered if appropriate. KEYTRUDA can also cause severe or life-threatening infusion-related reactions. Based on its mechanism of action, KEYTRUDA can cause fetal harm when administered to a pregnant woman. For more information, see "Selected

This indication was also based on data from KEYNOTE-180, a multicenter, non-randomized, open-label trial that enrolled 121 patients with locally advanced or metastatic esophageal cancer who progressed on or after at least two prior systemic treatments for advanced disease. With the exception of the number of prior lines of treatment, the eligibility criteria were similar to and the dosage regimen identical to KEYNOTE-181.