

BioSpect

the business of Bio

24 years of leveraging the business of Bio
www.biospectrumindia.com

How Precision Chemistry in Nobel-2026 can power pharma exports and national innovation

21 October 2026 | Views | By Professor Vidya Avasare, Department of Chemistry, Ashoka University

India can strengthen sovereign supply chains, boost high-value exports, and secure its place at the forefront of global pharmaceutical innovation



Asymmetric synthesis is a mature field in the construction of small to complex organic molecules. Why, then, was the 2026 Nobel Prize in Chemistry awarded to Henri B. Kagan and Kenso Soai for their discoveries regarding non-linear effects and asymmetric autocatalysis? How does it bring a fresh perspective to asymmetric synthesis and boost innovation in pharma industries?

Their discovery stems from nature's unmatched selectivity, observed in amino acids, carbohydrates, and complex natural products. For example, while the anticancer agent Taxol and the complex alkaloid strychnine mathematically possess 2,048 and 64 theoretical stereoisomers, nature exclusively synthesizes a single enantiomer without using complex transition metal catalysts or intricate ligands. How does nature yield only one mirror-image/optical isomer while suppressing the other? This breakthrough is at the core of the 2026 Nobel Prize.

Although Soai published his pioneering *Nature* paper on asymmetric autocatalysis nearly 30 years ago in 1995, the concept stems from Frank's 1953 model showing how mutual inhibition and autocatalysis amplify tiny stochastic fluctuations into a single isomer. By utilising an initial product with as little as 2% enantiomeric excess, the system self-amplifies throughout the reaction, completely eliminating the need for external chiral ligands, organocatalysts, or transition metal complexes.

To understand the value of their work, we must look back at history. The tragedy of thalidomide highlighted the critical importance of stereochemistry in drug design: while the (R)-enantiomer acted as a safe sedative, the (S)-enantiomer proved to be severely teratogenic, leading to severe birth defects. This disaster transformed pharmaceutical standards, making it mandatory for the industry to synthesise and evaluate optically pure compounds ensuring that only the single, biologically active enantiomer is delivered rather than a racemic mixture. Subsequent breakthroughs in asymmetric catalysis have dramatically impacted drug manufacturing by enabling single-enantiomer production.

However, as shown in **Table 1**, comparing these paradigm-shifting discoveries reveals unique pros, cons, and trade-offs across transition metal catalysis, organocatalysis, and non-linear asymmetric autocatalysis. Although used commonly and reliably, traditional chiral catalysts often face industrial limitations due to the high cost of synthesizing complex chiral ligands or noble metal complexes. Beyond expense, catalyst toxicity poses a major risk of heavy metal contamination in final drugs.

Furthermore, most homogeneous catalysts are difficult to recycle meaning they are synthesised, used once, and discarded as toxic waste. This exact industrial bottleneck is where the groundbreaking work of Henri Kagan and Kenso Soai opens an ambitious new frontier. By uncovering non-linear effects and asymmetric autocatalysis, their work demonstrates how exceptionally high enantiomeric purity can be generated while drastically minimizing unwanted side products and waste, much like nature does.

Honest reflection reminds us that a Nobel Prize does not transform an industry overnight. At many instances, such awards do not deliver immediate, off-the-shelf factory blueprints. Instead, their true power lies in breaking scientific boundaries, sparking new ideas, and establishing the beginning of a new technological era. While non-linear autocatalysis is currently limited to specific reactions, there is immense scope to expand these concepts to synthesize complex, pharmaceutically significant compounds at scale.

For India, this is a pivotal moment to capture. A large volume of Indian chiral API production still relies on traditional resolution techniques, which are energy-intensive, costly, and generate substantial chemical waste. While Europe, Japan, and the US hold majority ownership over patented chiral ligands and noble transition-metal catalysts, Indian manufacturers have historically depended on importing expensive chiral synthons or employing resolution steps that discard the undesired isomer as waste. This structural gap is precisely why adopting breakthrough concepts like non-linear effects and asymmetric autocatalysis, the Kagan-Soai principles is critical.

By reducing or eliminating reliance on costly, single-use metal catalysts and imported chiral ligands, Indian API manufacturers can drastically lower production costs, minimise environmental waste, and expand their global market share from ~15% toward leadership in high-value chiral drug discovery and CDMO exports.

Unlocking this "India Advantage" requires treating this scientific shift as a long-term strategic investment rooted in deep industry-academia collaboration. By bringing these precise chemistry techniques from fundamental theory down to the root level of our manufacturing plants, India can strengthen sovereign supply chains, boost high-value exports, and secure its place at the forefront of global pharmaceutical innovation. In my opinion, this marks the true beginning of non-linear autocatalysis in industrial applications, we must take a lion's leap to expand this idea to elevate our global contribution.

Table 1: Comparison of Nobel Prize-Winning Asymmetric Catalysis

Parameters	Asymmetric Transition Metal Catalysis	Asymmetric Organocatalysis	Asymmetric Autocatalysis
Recognition	2001 Nobel Prize	2021 Nobel Prize	2026 Nobel Prize
Primary Catalyst Source	Transition metals coordinated with complex chiral ligands	Small chiral organic molecules directly from nature like proline without metal ions	The reaction product itself is a catalyst
Synthesis & Material Cost	Very High due to precious metals and complex ligands	Low to Moderate: Readily available from natural chiral pool molecules	Almost Zero: No external chiral ligand
Purity Needed	High (>98% ee)	High (>95% ee)	Low (<2% ee or traces)
Catalyst Loading	Low to moderate (typically 0.1 to 10 mol%)	High (often 10 to 20 mol%)	Negligible / Traces
Productivity (TOF / TON)	Very High: Fast reaction kinetics, high turnover frequencies (TOF) and turnover numbers (TON)	Low to Moderate: Slower reaction speeds; higher reaction times required	Exponential Rates: Characterized by an induction phase followed by exponential acceleration as catalyst concentration increases
Recyclability	Difficult & Expensive	Moderate to Difficult	Self-Resolving: The catalyst is released as a byproduct

Toxicity & Heavy Metal Residue	High Concern: Severe risk of toxic heavy metal contamination in final Active Pharmaceutical	Low Toxicity: Environmentally benign; no heavy metals involved	Low to Variable: Depends on main reagents/substrates involved; avoiding toxic precious metal contamination
Environmental Impact & Waste (E-factor)	High: Heavy metal extraction, ligand synthesis waste, and single-use disposal create high environmental burdens	Low: Green chemistry credentials; biodegradable and non-metal reagents	Very Low: Eliminates dedicated manufacturing streams and minimizes toxic waste generation
Substrate Scope & Applicability	Extremely Broad: Handles a vast array of transformations (hydrogenations, cross-couplings, C–H activations)	Broad: Excellent for aldol, Mannich, conjugate additions, and organo-reductions	Currently Specialised: Scope restricted to specific self-amp systems (e.g., dialkylzinc additions to aldehydes)
Sensitivity to Air & Moisture	Extremely high	Low	Moderate to High: Depends on organometallic reagents used in specific reaction manifold

Professor Vidya Avasare, Department of Chemistry, Ashoka University

(The views expressed in this article are the author's personal opinions and do not necessarily reflect those of Ashoka University)